



15-0-3

Turf King Fertilizer

with 0.20% Prodiamine

For pre-emergence control of grass and broadleaf weeds in lawns and ornamental turf (excluding golf course putting greens)

Landscape ornamentals, established perennials and wildflower plantings

Stops many weed grasses before they grow - Prevents crabgrass, annual bluegrass, foxtail

ACTIVE INGREDIENT:

Prodiamine: (CAS # 29091-21-2)..... 0.20%

OTHER INGREDIENTS:..... 99.80%

TOTAL:..... 100.00%

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for further precautions and First Aid Information below

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

FIRST AID

IF IN EYES:	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	- Take off contaminated clothing. - Rinse skin immediately with plenty of water for 15-20 minutes. - Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	- Call a poison control center or physician immediately for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by the poison control center or physician. - Do not give anything by mouth to an unconscious person.
IF INHALED:	- Move person to fresh air. - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. - Call a poison control center or doctor for further treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall® International (866) 897-8050 for emergency medical treatment information.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

Distributed and Guaranteed by Andre & Son
17150 State Route 706 • Montrose, PA 18801 • 570-278-1131

GUARANTEED ANALYSIS

Total Nitrogen (N) 15.00%
11.25% Urea Nitrogen
3.75.% CSRUN

Potash (K₂O) 3.00%

Nutrients derived from:

XCU, Urea, Muriate of Potash.

Information concerning the raw materials composing this product can be obtained by writing to the Guarantor listed on this label, and referring to the batch number found on this container.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

This product is a pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established lawns and ornamental turf (excluding golf course putting greens), landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development. When used according to directions, it will control; barnyardgrass; bluegrass; annual (Poa annua); carpetweed; chickweed, common; chickweed, mouseear (from seed); crabgrass (large, smooth); crowfootgrass; cupgrass, woolly; foxtails, annual; goosegrass1; henbit; itchgrass; johnsongrass (from seed); junglerice; knotweed; kochia; lambsquarter, common; lovegrass; panicum (Texas, fall, browntop); pigweed; purslane, common; pusley, Florida; rescuegrass3, shepherd's purse2, signalgrass, broadleaf; speedwell, Persian; sprangletop; spurge, prostrate; witchgrass; woodsorrel, yellow (from seed).

1In many areas a single application of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 7.6 to 11.5 pounds/1,000 square feet (330 to 500 pounds/acre), followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Minimum and Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

2For this weed, apply in late summer, fall or winter prior to germination.

3Suppression only.

NET WT 50 lb (22.7 kg)

EPA Reg. No. 53883-173-47956 • EPA Est. No. 47956-PA-001

WHEN TO APPLY TO ESTABLISHED TURF:

This product must be applied prior to germination of weed seeds, as it will not control established weeds.

THIS MAP SHOWS APPROXIMATE CRABGRASS GERMINATION DATES



APPLICATION DIRECTIONS:

Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1-2 inches) incorporation, prior to weed seed germination and within 14 days application.

USE PRECAUTIONS:

The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may thin emerged annual bluegrass and newly overseeded grasses. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil). Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (Poa annua) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

RATES OF APPLICATION:

This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

MINIMUM AND MAXIMUM APPLICATION RATES (by Turf Grass Species)		
Turf Species	Minimum Rate	Maximum Annual Rate ¹
Bahiagrass, Bermudagrass ² Centipedegrass, Zoysiagrass, Seashore Paspalum, St. Augustinegrass Tall Fescue (including turf-Type)	4 lbs/1,000 sq. ft. (174 lbs/acre)	17.2 lbs/1,000 sq. ft. per year (750 lbs/acre per year)
Buffalograss Kentucky Bluegrass Perennial Ryegrass	4 lbs/1,000 sq. ft. (174 lbs/acre)	11.5 lbs/1,000 sq. ft. per year (500 lbs/acre per year)
Creeping Red Fescue	4 lbs/1,000 sq. ft. (174 lbs/acre)	8.5 lbs/1,000 sq. ft. per year (370 lbs/acre per year)
Creeping Bentgrass	4 lbs/1,000 sq. ft. (174 lbs/acre)	7.6 lbs/1,000 sq. ft. per year (330 lbs/acre per year)

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 6 pounds/1,000 square feet (260 pounds/acre). Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control

DIRECTION FOR USE (continued)

WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results. (See map)

USE RATE	MONTHS BEFORE OVERSEEDING		
	North	Transition	South
4 pounds/1,000 sq. ft. (174 pounds/acre)	4	4	4
7.6 pounds/1,000 sq. ft. (330 pounds/acre)	5	4	4
8.5 pounds/1,000 sq. ft. (370 pounds/acre)	6	5	5
9.2 pounds/1,000 sq. ft. (400 pounds/acre)		6	6
11.5 pounds/1,000 sq. ft. (500 pounds/acre)		7	7
13 pounds/1,000 sq. ft. (570 pounds/acre)			9
14.9 pounds/1,000 sq. ft. (650 pounds/acre)			10
17.2 pounds/1,000 sq. ft. (750 pounds/acre)			12

LANDSCAPE ORNAMENTALS (including established perennials and wildflower plantings):

This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 7.6 to 17.2 pounds/1,000 square feet (330 to 750 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 17.2 pounds/1,000 square feet (750 pounds/acre) per year.

SPREADER SETTINGS to apply at the listed pounds per 1,000 square feet

SR2000	4 lbs	6 lbs	7.6 lbs	8.5 lbs	11.5 lbs	17.2 lbs
	M	O	Q	R	S 1/2	

(May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product)

APPLICATION TIMING AND INFORMATION

This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Do not rinse unless required for recycling. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

CONDITIONS OF SALE AND WARRANTY

The DIRECTIONS FOR USE of this product are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop, injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of Andre & Son or the Seller. All such risks shall be assumed by the Buyer.

Andre & Son warrants that this product conforms to the chemical description on the label, and is reasonably fit for the purposes set forth in the DIRECTIONS FOR USE, when it is used in accordance with such directions, subject to the inherent risks mentioned above.

Andre & Son NEITHER MAKES NOR INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. THIS WARRANTY EXTENDS TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNING, OR CAUTIONS. BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION TO REPLACEMENT OF, OR THE REPAYMENT OF THE PURCHASE PRICE FOR, THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Andre & Son AND THE SELLER OFFER THIS PRODUCT, AND THE BUYER AND USER ACCEPT IT, SUBJECT TO THE FOREGOING CONDITIONS OF SALE AND WARRANTY.

TOLERANT ORNAMENTAL SPECIES:

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME	COMMON NAME	SCIENTIFIC NAME	COMMON NAME
Abies spp.	Fir species** (Balsam, Frasier, Noble, etc.)	Hedera helix	English Ivy	P. taeda	Loblolly Pine
Abelia grandiflora	Abelia: Sherwood	Hamamelis virginica	Shagbark	P. thunbergiana	Japanese Black Pine
Acer palmatum	Japanese Maple	Hemerocallis spp.	Daylily: Aztec Gold, Stella de Oro, Tender Love	P. virginiana	Virginia Pine
A. platanoides	Norway Maple	Hibiscus	Rose of Sharon**	Pistachio spp.	Pistachio*
Achillea spp.	Yarrow: King Edward	Hibiscus spp.	Mallow: Disco Belle White	Pittosporum rhombifolium	Queensland Pittosporum
Actinidia chinensis	Kiwi*	Hibiscus rosa-sinensis	Chinese Hibiscus	P. tobira	Japanese Pittosporum
Agapanthus africanus	Lily-of-the-Nile (African Lily)	Hosta plantaginea	Hosta, Plantain Lily (Fragrant)	Podocarpus macrophyllus	Japanese Yew
Agapanthus orientalis		H. sieboldiana	Hosta, "Seersucker"	Prunus laurocerasus	English Laurel
Akebia quintata	Five-leaf or Chocolate Vine	Houttuynia cordata var. variegata		Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Allium cernuum	Lady's Leek, Nodding Onion	Hydrangea macrophylla	Bigleaf Hydrangea	Pseudotsuga menziesii	Douglas Fir**
Anemone hybrida	Japanese Anemone	Ilex Bernyi	Holly	Pyraecanthia coccinea	Firethorn (Scarlet)
Aquilegia spp.	Aquilegia: Red and Gold	Ilex cornuta	Chinese Holly	P. fortuneana	Firethorn
Arctostaphylos densiflora	Vine Hill Manzanita	I. crenata	Japanese Holly	P. koidzumii	Firethorn
Arctotheca calendula	Cape Weed	I. opaca	American Holly	Pyrus spp.	Bradford Pear spp.
Artemisia spp.	Wormwood; Silver Mound, Castle	I. vomitoria	Yaupon Holly, Schillings	Quercus shumardii	Oak, Shumard's Red
Aster spp.	Aster: Bonnie Blue, Purple Dome	Inula ensifolia		Quercus spp.	Oak species
Aster X frikartii		Iris ensata	Sword-Leaved Iris; Jodlesong	Raphiolepis indica	Indian Hawthorne
Athrium filiz-femina	Lady Fern; Fern Lady	Iris sibirica	Siberian Iris; Cabernet	Raphiolepis umbellata	Yedda Hawthorne
Aucuba japonica	Japanese Aucuba	Iris spp.	Iris species**	Rhododendron	'Coral Bells,' 'Formosa,' 'Hino-crimson,' 'PJM'
Begonia spp.	Fibrous Begonia: Hardy Grandis	Jasminium nudiflorum	Winter Jasmine	(including Azalea)	'Delaware Valley White,' 'Roseum Elegans,' 'Flame Creeper', 'Girard Crimson,' 'Wakeiebusu 'White Gumbo', 'George L. Tabor'
Berberis gladiolifera	Barberry	Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii	Rosa banksiae	Lady Bank's Rose
B. julianae	Wintergreen Barberry	J. conferta	Shore Juniper; Blue Pacific	Rosmarinus officinalis	Rosemary*
B. mentorensis	Mentor Barberry	J. davurica	Parsoni	Rudbeckia spp.	Black-Eyed Susan; Goldsturm
B. thunbergii	Japanese Barberry	J. thunzialis	Creeping Juniper	Santolina virens	Santolina
B. verruculosa	Warty Barberry	Juglans spp.	Walnut*	Saxifraga spp.	Saxifrage; Purple Dome
Bergenia cordifolia		Justicia brandegeana	Shrimp Plant	Scabiosa spp.	Pincushion Flower
Boltonia asteroides	Snowbank	Lagerstromia indica	Crape Myrtle	Sedum album	Stoncrop
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red	Lantana montevidensis	Weeping Lantana	Sedum caudicula	Stoncrop; Lidakense
Buxus microphylla	Japanese Boxwood	Lavender spp.	Edelweiss	Sedum dasyphyllum	Stoncrop
Callistemon viminalis	Weeping Bottlebrush	Leonopodium alpinum	Amur Privet	Sedum spurium	Stoncrop; Dragon's Blood
Calluna vulgaris	Tussock Bellflower, (White Clips)	Ligustrum amurense	Chinese Privet; Variegata	Spiraea bumalda	Spiraea; Anthony Waterer
Campanula carpatica	Trumpet Flower, Madame Galen	Ligustrum sinense	Japanese Privet	Syzygium paniculatum	Australian Brushcherry
Campis X tagliabuana	Hottentot Fig (Ice Plant)	L. japonicum	Glossy Privet (Waxleaf)	Taxus cuspidata	Japanese Yew
Cassia artemisoides	Feathery Cassia	L. lucidum	Lily; Jazz	T. media	Yew
Ceanothus rigidus	Wild Lilac	Lilium spp.	Big Blue Lily, Silver Mound, Evergreen Giant	Teucrium spp.	Germander
Ceratostigma plumbagonoides	False Cypress	Liriope muscari	Liriope, Variegated	Thalictrum dipteracarpum	Meadow Rue
Chamaecyparis pisifera	Cleyera	Liriope muscari var. variegata	Liriope, Creeping	Thuja occidentalis	American Arborvitae
Cleyera japonica	Citrus species*	Liriope spicata	Lobelia Cardinalis	Trachelospermum asiaticum	Star Jasmine
Citrus spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam	Lobelia cardinalis	Japanese Honeysuckle	Tsuga canadensis	Canada Hemlock
Coreopsis spp.	Flowering Dogwood	Lonicera japonica	Tatarian Honeysuckle	Tulipa spp.	Tulip species
Cornus florida	American Dogwood	L. tatarica	Burgundy	Veronica spp.	Veronica, Speedwell; Sunny Border
C. stolonifera	Pampas Grass	Lorapetalum chinense	Magnolia species**	Viburnum japonicum	Sweet Viburnum
Cortaderia selloana	Cranberry Cotoneaster	Lythrum spp.	Ice Plant	V. odoratissimum	Japanese Snowball
Cotoneaster apiculatus	Cotoneaster	Magnolia spp.	Crabapple*	V. plicatum	Canary Island Viburnum
C. buxifolius	Bearberry Cotoneaster	Maleophora luteola	Yaku Jima, Silberfeder	V. suspensum	Arrowwood Viburnum
C. dammeri	Rockspray Cotoneaster	Malus spp.	Heavenly Bamboo	V. tinus	Laurustinus
C. microphyllus	Hawthorne	Miscanthus sinensis	Narcissus species**	V. trilobium	Cranberry Bush
Cretaeus spp.	Lucifer	Nandina domestica	Oleander	V. wrightii	Leatherleaf Viburnum
Crocossmia spp.	Italian Cypress	Narcissus spp.	Olea europaea	Vinca major	Greater Periwinkle, Vinca
Cupressus sempervirens	White Trailing Ice Plant	Nerium oleander	Ophiopogon japonicus	Vinca minor	Common Periwinkle, Vinca
Delosperma alba	Cooperi Pink	Olea europaea	Omanthus heterophyllus	Vitis spp.	Grape*
Delosperma spp.	Larkspur: Blue Elf	Ophiopogon japonicus	Osteospermum fruticosum	Weigela florida	Old Fashioned Weigela
Delphinium spp.	Dianthus: Maiden Pinks "Zing"	Omanthus heterophyllus	Oxydendrum arboreum	Yucca aloifolia	Spanish Bayonet, Yucca
Dianthus deltoideus	Cheddar Pink	Osteospermum fruticosum	Pennisetum setaceum	Y. filametosia	Adam's Needle, Yucca
D. gratianopolitanus	Hop Bush	Oxydendrum arboreum	Perovskia atriplicifolia		
Dodonea viscosa	Coneflower, Purple; Magnus	Paeonia suffruticosa	Persea americana		
Echinacea purpurea	Silverberry	Pennisetum setaceum	Photinia Fraseri		
Elaeagnus pungens	Wintercreeper	Perovskia atriplicifolia	Physotelia virginiana		
Euonymus fortunei	Evergreen Euonymus	Photinia Fraseri	Picea spp.		
E. japonica	Spreading Euonymus	Physotelia virginiana	Spruce species** (Colorado Blue, Norway, etc.)		
E. kiautschovica	Japanese Aralia	Picea spp.	Lily-of-the-Valley Bush		
Fatsia japonica	Border Forsythia	Pieris japonica	Calabrian Pine		
Forsythia intermedia	Weeping Forsythia	Pinus brutia	Canary Island Pine		
Forsythia suspensa	Greenstem Forsythia	P. canariensis	Slash Pine		
F. viridissima	Gaillardia, Blanket Flower: "Goblin"	P. Elliottii	Aleppo Pine		
Gaillardia spp.	Gardenia, Cape-Jasmine	P. halepensis	Austrian Pine		
Gardenia jasminoides	Gentian	P. nigra	Longleaf Pine		
Gaura spp.	Cranesbill	P. palustris	Monterey Pine		
Gentiana dahurica	Gladiolus spp.**	P. radiata	Eastern White Pine		
Geranium cinereum	Baby's Breath	P. strobilus	Scotch Pine		
Gypophylla repens		P. sylvestris			

*Do not use on food producing trees, vines or plants.

**Not for use on container grown plants.